Review Article

Treatment of traumatic rib fractures: an overview of current evidence and future perspectives

Aimée J. P. M. Franssen[1#^], Jean H. T. Daemen[1#^], Julien A. Luyten[1^], Berry Meesters[2^], Annette M. Pijnenburg[2], Kostan W. Reisinger[2], Raoul van Vugt[2^], Karel W. E. Hulsewé[1^], Yvonne L. J. Vissers[1^], Erik R. de Loos[1,2^]

[1]Division of General Thoracic Surgery, Department of Surgery, Zuyderland Medical Centre, Heerlen, the Netherlands; [2]Division of Trauma Surgery, Department of Surgery, Zuyderland Medical Centre, Heerlen, the Netherlands

Contributions: (I) Conception and design: AJPM Franssen, JHT Daemen, ER de Loos; (II) Administrative support: AJPM Franssen, JHT Daemen; (III) Provision of study materials or patients: B Meesters, AM Pijnenburg, KW Reisinger, R van Vugt, KWE Hulsewé, YLJ Vissers, ER de Loos; (IV) Collection and assembly of data: AJPM Franssen, JHT Daemen, JA Luyten; (V) Data analysis and interpretation: AJPM Franssen, JHT Daemen, ER de Loos; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

[#]These authors contributed equally to this work.

Correspondence to: Erik R. de Loos, MD, PhD. Division of General Thoracic Surgery, Department of Surgery, Zuyderland Medical Centre, Henri Dunantstraat 5, 6419PC Heerlen, the Netherlands; Division of Trauma Surgery, Department of Surgery, Zuyderland Medical Centre, Henri Dunantstraat 5, 6419PC Heerlen, the Netherlands. Email: e.deloos@zuyderland.nl.

Abstract: Operative management of rib fractures has gained significant popularity over the last years, however, it remains a controversial topic, due to the substantial heterogeneity among rib fracture patients with considerable differences in epidemiology. Hence, the present narrative review aims to provide an overview of the treatment and (long-term) outcomes of rib fractures, with an emphasis on the surgical treatment. Nowadays, computed tomography (CT) has been shown to be most practical and sensitive for detecting rib fractures, of which up to 50% is missed on other imaging modalities. Non-operative treatment by patient-tailored multimodal pain management remains the cornerstone. Still, in the presence of—amidst others—chest wall instability or displaced fractures with physiologic derangements, operative treatment is indicated and should be performed within 72 hours after injury. Here, traumatic brain injury (TBI) and pulmonary contusion are no strict contra-indications, while plate osteosynthesis is considered the standard mode for surgical stabilization. To date, surgical stabilization of rib fractures (SSRF) only benefits selected groups of patients, awaiting results of ongoing studies. Future directions may include the sole use of percutaneous cryoablation of the intercostal nerves as part of conservative management, as well as the application of three-dimensional (3D) printing and use of bio-absorbable materials in the surgical treatment of rib fractures.

Keywords: Rib fractures; thoracic injuries; non-operative management; surgical stabilization of rib fractures (SSRF)

Submitted Dec 02, 2023. Accepted for publication Jun 26, 2024. Published online Jul 26, 2024.
doi: 10.21037/jtd-23-1832
View this article at: https://dx.doi.org/10.21037/jtd-23-1832

^ ORCID: Aimée J. P. M. Franssen, 0000-0001-6399-1663; Jean H. T. Daemen, 0000-0002-4878-3951; Julien A. Luyten, 0000-0003-2350-1521; Berry Meesters, 0009-0002-6069-1627; Raoul van Vugt, 0000-0002-3293-7223; Karel W. E. Hulsewé, 0000-0001-8131-1895; Yvonne L. J. Vissers, 0000-0002-2890-8390; Erik R. de Loos, 0000-0001-6313-2658.

© Journal of Thoracic Disease. All rights reserved.

SHOUT
SHARE
IT

Introduction

Thoracic injury is a common and potentially devastating component of acute trauma. The majority—90%—of thoracic injuries result from blunt trauma, accounting for 14% of all blunt traumas. Furthermore, thoracic trauma is associated with significant morbidity and mortality, estimated to be accounting for up to a quarter of early trauma-related mortality (1). Regarding thoracic trauma-related specific injuries, rib fractures are considered the most prevalent, with a large variety in severity, ranging from simple isolated fractures to multiple and multifragmentary fractures resulting in flail chest, for example in polytrauma patients. Despite the generally urgent nature of thoracic trauma, after quick diagnosis most injuries can be mitigated by non-operative management or later definitive surgical treatment. Operative management of rib fractures has gained significant popularity over the last years, however, it remains a controversial topic. The substantial heterogeneity among rib fracture patients with considerable differences in epidemiology (2) perhaps prevents a one-size-fits-all solution to rib fracture management. Hence, the present narrative review aims to provide an overview of the different treatment strategies and (long-term) outcomes of rib fractures, with an emphasis on the surgical treatment.

Initial management and diagnosis of thoracic injuries

Initial resuscitation and management of acute chest trauma patients is based upon the principles of Advanced Trauma Life Support (3). To evaluate the presence of acute life-threatening thoracic injury at an early stage, an important adjunct in primary survey is plain chest radiography. Although acquired in all trauma patients, chest radiographs are associated with poor diagnostic accuracy (4,5). Approximately 50% of rib fractures are missed on routine chest radiographs (5,6). Oblique and special rib views can increase sensitivity but are not routinely obtained. Chest ultrasound can serve as an alternative to detect thoracic injuries such as traumatic hemopneumothorax and rib fractures. Although ultrasonography is associated with even poorer diagnostic accuracy (7) to detect a traumatic pneumothorax, it has shown superior results in the detection of rib fractures compared to chest radiographs (8). Still, current guidelines do state chest radiographs as the most appropriate primary imaging modality in adults with suspected rib injuries (9). This may be due to the variability in patient habitus and fracture sites (more difficult to visualize in the subscapular and infraclavicular portion) (10), as well as the increased imaging time associated with ultrasound (11). Nuclear imaging could in extremely rare circumstances be used to detect occult rib fractures but should definitely have no place in acute trauma care (12). Computed tomography (CT) has been shown to be most practical and sensitive for detecting rib fractures (6), hence often obtained subsequent to simple radiographs and considered gold standard. The importance to detect rib fractures is underlined by the fact that the pulmonary morbidity and mortality directly increases with each additional rib fracture, approaching a mortality of 40% if more than six ribs are fractured (13), prompting more aggressive management. CT moreover allows for a comprehensive evaluation of the chest, identifying injuries such as pulmonary contusion, a diaphragmatic rupture and entrapped lung parenchyma. In addition, three-dimensional (3D) reconstructions are useful for pre-operative surgical planning.

Non-operative management

In the absence of a flail chest or chest wall deformity (including e.g., severely displaced ribs), non-operative management remains the standard treatment of thoracic injuries at most institutions.

Healing thus usually occurs spontaneously and is beneficial if it avoids potential surgery-related complications. On the other hand, insufficient bone healing may result from rib fractures that are not surgically repositioned and stabilized, resulting in the development of pseudarthrosis and chronic pain syndromes.

Notwithstanding, spontaneous bone healing is associated with substantial pain and discomfort. Hence, the mainstay of conservative management is adequate pain control to prevent secondary pulmonary complications resulting from inadequate ventilation due to painful respiratory movements. This is best achieved by a staged protocol. It is advised to directly start analgesia at the emergency department by a combination of acetaminophen, non-steroid anti-inflammatory drugs (NSAIDs) or metamizole in the presence of contra-indications, and opioids. Repeated evaluation of pain scores must be performed, both in rest and during exercise or coughing. Inadequate pain control with numeric rating scores of >4 should initiate the placement of a thoracic epidural catheter. A systematic review by Peek and colleagues found that epidural analgesia provides better pain relief than other modalities [including intravenous analgesia

© Journal of Thoracic Disease. All rights reserved.

and (loco-)regional blocks], though no differences were observed for the length of intensive care unit (ICU) stay and mechanical ventilation and pulmonary complications (14). Another study by Jensen and colleagues predicted a 97% reduction in mortality with thoracic epidural analgesia among patients with one or multiple rib fractures (15), while others particularly advocated for epidural use with more than four rib fractures due to the associated mortality risk reduction (13). In the presence of contra-indications, such as the use of specific anticoagulants, one can refer to a thoracic erector spinae plane block or paravertebral block, which may be of equivalent effect (16,17). Though, it must also be mentioned that different studies question the efficacy and safety of epidural catheters (18,19).

Persistent high pain scores despite adequate non-operative measures should reroute patients towards operative treatment.

Without going into the multitude of individual methods of pain relief, choosing the optimal technique may not always be straightforward, and thus treatment should always take place in cooperation with a pain specialist, respiratory physiotherapist, be multimodal, patient-specific and tailored to local expertise and experiences (20). In addition, one should keep in mind that despite trying to avoid surgery and the associated risks, conservative treatment can also have extremely rare devastating consequences, such as epidural hematoma or abscess formation following epidural catheterization (21).

Over the years, different scoring systems have been developed to identify patients at risk for adverse outcomes. These include the SCARF (22) and BATTLE score (23) and may guide treatment decisions. Predictors of adverse outcomes encompassed in these scores are age, number of rib fractures, presence of chronic lung disease, anticoagulant use, oxygen saturation levels, increased respiratory rate, high pain scores and inadequate coughing, prompting more aggressive management. The thoracic trauma severity score (TTSS) has been shown to be superior specifically in predicting respiratory complications in isolated rib fractures with pulmonary contusion and the rib fracture score (RFS) was proven most useful in patients without pulmonary contusion compared to the abbreviated injury scale (AIS), chest trauma score (CTS) and RibScore (24).

Operative management

Over the last two to three decades, surgical stabilization of rib fractures (SSRF) has gained popularity compared to conservative treatment options, especially for flail chest (25,26). In general, the assumption that SSRF could alleviate respiratory problems remains the main driver for operative management, as was affirmed by Dutch health care providers using the measurement instrument for determinants of innovations (MIDI) questionnaire, which was used to identify barriers and facilitators of the implementation of SSRF (27). Although the skepticism towards SSRF is largest for non-flail chest patients, different studies suggest suboptimal quality of life and increased morbidity even after isolated rib fractures (28,29), creating opportunities for surgical fixation in selected patients.

The Chest Wall Injury Society (CWIS), represented by a variety of surgical and non-surgical specialists, seeks optimal outcomes for patients with both non-operative and operative management of rib fracture. Being leaders in rib fracture research, CWIS members have reported an evidence-based consensus guideline for indications, timing, surgical approach, and fixation mode of SSRF (30).

Indications

CWIS set distinct recommendations on the indications to perform SSRF for both ventilated and non-ventilated patients. Although the indications slightly differ between both groups, SSRF is generally indicated in the presence of chest wall instability (e.g., flail chest, clinical instability, and paradoxal chest motion), three or more displaced rib fractures (equal to or more than 50% of the rib width in costae 3 to 10) with multiple pulmonary physiologic derangements (e.g., high respiratory rate and pain scores despite adequate multimodal pain management), or failure to wean from ventilation (31). The society also outlined contraindications for SSRF, both absolute and relative. Absolute contraindications include severe traumatic brain injury (TBI) and acute myocardial infarction, while less than severe TBI and spinal injuries are more relative. It has to be mentioned though that since the publication of CWIS guidelines, new evidence has come to light about the benefits of SSRF for patients with multiple rib fractures and simultaneous moderate to severe TBI, despite TBI being a driver for acute respiratory outcomes (32,33). As such, it has been shown that there is no SSRF-related complication risk, and SSRF treatment resulted in a lower 30-day mortality in patients with severe TBI compared to conventional non-operative treatment (34).

© Journal of Thoracic Disease. All rights reserved.

Optimal timing

Timing of surgical rib fracture repair is essential and should consider inflammation and callus formation. Furthermore, fractures are known to increasingly displace with time, setting hurdles for anatomic reconstruction. According to CWIS guidelines and a recent study by Prins and colleagues, SSRF should be performed as early as possible (i.e., within 24 hours) and ideally within 72 hours for non-ventilated patients regardless of the indication (31,35). When respiratory failure/ventilator dependence has complicated rib fractures in ventilated patients, SSRF should be performed within 72 hours for both flail as non-flail indications. It is recommended though that SSRF should be delayed in case of other higher priority injuries (e.g., spinal or vascular injuries, pulmonary contusion, or open abdomens), and hemodynamic instability which often necessitates emergency surgery. Especially pulmonary contusion has been considered a relative contra-indication as the risk of pulmonary morbidity is believed to be associated with the contusion itself instead of the rib fractures (36). Recently, a study by Lagazzi and colleagues did show that regardless the severity of the pulmonary contusion, early implementation of SSRF (i.e., <72 hours) is advised, as it was associated with a decrease in hospital and ICU stay, and more importantly lower rates of unplanned intubation or tracheostomy compared to late SSRF (i.e., >72 hours) (37). Following up on this study, Van Wijck and colleagues showed that there is no association between pulmonary contusion and worse SSRF outcomes in patients with three or more displaced rib fractures or flail segment (33). They even showed that SSRF is associated with superior outcomes in patients with mild to moderate pulmonary contusion. Nevertheless, patients experiencing severe pulmonary contusion should be considered for SSRF on an individual basis as outcomes did not differ between surgical and non-operative management. Regardless of the indications and timing for SSRF, it is recognized that the choice of treatment and timing remains partially up for debate because each patient and associated trauma is unique, as are the subjective skills of the surgeon and resource options.

Surgical approach

Conventionally, the SSRF era started with the use of large incisions obtaining full exposure, however the corresponding need to muscle and nerve stripping and potential damage to blood vessels brought along several complications (e.g., post-operative scarring, functional limitations, numbness, wound infection). Currently, SSRF may be assisted by video-assisted thoracoscopic surgery (VATS), offering a wide visualization of the entire chest wall, although not considered mandatory. Addition of VATS has shown to be effective and safe and allows for the identification and repair of additional pulmonary or mediastinal (pathologic) conditions as well as the simultaneous evacuation of for example a hemothorax (25,38,39). However, it has been shown that addition of uniportal VATS to SSRF on a routine basis for severely displaced rib fractures or flail chest does not identify a significantly greater amount of occult intrathoracic injuries compared to using uniportal VATS only on selected cases (40). Notwithstanding, VATS has the function of identifying clinically relevant rib fractures, as not every single fracture requires stabilization through osteosynthesis; combined with intraoperative palpation of the thorax, clinically relevant fractures can be specifically identified and the most atraumatic approach to the corresponding ribs can subsequently be determined.

In theory, VATS-assisted SSRF has also the advantage of video-guided placement of loco-regional anesthesia, which is imperative to successful and proper treatment of rib fractures, as well as chest tube placement. In addition, it can improve the visualization of rib fractures, allowing more accurate fracture identification. Moreover, it allows intrathoracic plate placement to the inner cortex of the rib potentially leading to better healing of the fracture and improved pain control compared to extrathoracic plate placement (41,42). One may even take intrathoracic placement one step further using uniportal instead of multiportal VATS. Although the data on this uniportal VATS SSRF approach is currently limited (43,44), its position is expected to be elucidated in the coming years.

While the use of VATS, either being used in addition to or as a complete procedure, is being gradually accepted and applied for surgical rib fixation, no comparing studies between VATS and thoracotomy for SSRF have been reported yet. On this account, no evidence-based conclusions can be drawn yet whether which approach is superior to the other.

Fixation mode

The ideal SSRF fixation product has a unique geometrical shape that respects the anatomical curvature of the rib in question (i.e., plasticity and biomechanical elasticity).

© Journal of Thoracic Disease. All rights reserved.

To restore lung volumes and respiratory function, it also requires sufficient strength and stiffness to reestablish chest wall stability. Furthermore, it has good histocompatibility, causes no adverse reaction, and does not affect surrounding tissues. While there is currently no product that perfectly covers all these aspects, CWIS does lay out specific recommendations for the mode of fixation (31). When multiple fractures are identified, all fracture lines should be stabilized wherever possible (45). Since there is no clinical evidence available that suggests superiority of outer cortical plate fixation over intramedullary struts, both may be used to surgically fixate rib fractures. However, intramedullary struts are quite tricky to implant though and only have one anchor point, without distal fixation (46) and are generally considered inferior to cortical plate fixation. Nevertheless, cortical plate fixation is considered time consuming. Another option is to use nitinol plates/clips which apply the properties of shape memory alloys such as NiTi Rib plates and prevent the need for manual bending of the plate (47). When cooled down before application, the material is malleable, and only after heating up to body temperatures by placement onto the fracture site, the plates and clips undergo a phase transformation and return to their original shape.

While bicortical fixation is designed for ribs to be immobilized at a stiffness level optimal for bone healing, a unicortical approach can be used to prevent bicortical-induced theoretical risks of injuring intrathoracic organs, specifically pleural irritation or risk of pneumothorax by the prominence of the screw tip (48). Systems designed for a unicortical approach (such as the L1® Rib system from KLS Martin Group) place converging screws, aiming to match the biomechanical properties of a bicortical approach, including pull-out forces.

Other fixation options include Judet struts and Kirshner wires. The latter are used for transfixation where non-fractured ribs provide stabilization. Although the use of Kirshner wires is characterized for a reduction of operative time, it is associated with increased postoperative morbidity compared to Judet struts and cortical plate fixation (49). For these as well as other reasons (e.g., suboptimal fracture reduction and stability, as well as the risk for injury to surrounding structures), wires are hardly used at present.

Bioabsorbable prostheses for fracture fixation are not a new development, but are mostly utilized in other fields of surgery. These bioabsorbable and bioactive plates consists of poly-L-lactide and hydroxyapatite, the combination of which has osteoconductive properties that bond bones and thereby promotes bone healing. They especially provide strength and stiffness until the fractures are fully healed, and with their slow absorption rate mechanical stresses are slowly transferred to the bone, minimizing stress shielding and the need for surgical removal of foreign body materials. Such bioabsorbable mini plates and screws have been reported in the successful treatment of non-trauma patients who received chest wall reconstructions (50) as well as pectus excavatum patients (51). However, it was not until Mayberry and colleagues first showed good clinical use in surgical rib fracture stabilization in 2003 (52), that is has been recognized as a new revolutionary technique in the field of SSRF. Since then, only few studies have implemented bioabsorbable plates to further enhance bone healing of rib fractures (53-55). While stability issues remain with bioabsorbable plates, these internal fixation materials could be the main direction for future development.

Besides the different types of hardware that can be applied to surgically fixate rib fractures, one must always keep the basic principles of fracture management in mind and consider whether the additional use of bone grafts is necessary.

Outcomes

The question remains whether the enthusiasm for SSRF developed over the years is worth the cost. The overall risk of surgery- and implant-related complications has been reported in 10.3% of the cases, with wound infection in 2.2% and fracture-related infection in 1.3% (56). Risk factors for morbidity and mortality after SSRF, including age and number of fractured ribs, have however been reported inconsistently in the literature (57). Hence, identifying reliable predictors can have significant implications for the clinical outcomes of rib fracture patients, independent of treatment option. In 2012, a simple scoring system has been developed that may assist in the earlier implementation of treatment strategies (58).

While the short-term clinical patient outcomes after SSRF are clearly described in the literature (59-62), with less pain and respiratory-related complications/readmissions, higher rates of home discharge, a decrease in ventilator time and tracheostomy and in-hospital costs, more studies are now investigating long-term benefits of SSRF. Patients receiving SSRF for their rib fractures can return to their usual life or same work as in their pre-injury state with no or few limitations besides from implant-

© Journal of Thoracic Disease. All rights reserved.

related irritation (63-65). Nevertheless, most studies exclude the patient subgroup of 80 years and older, while it has been shown that specifically in this subgroup SSRF is independently associated with a decreased mortality (66). Thus, specific patient selection is required with more randomized controlled trials. So far, only a few RCTs on surgical versus non-operative treatment for severe chest wall injury have been reported in the literature. While these studies have shown less restrictive impairment of pulmonary function (67), a lower risk of complications as well as less ventilation requirement, intensive care stay, and thus medical resource use after surgical treatment (68,69), most patients with severe chest wall injury still experience moderate to extreme pain as well as impaired physical activity at 1 month after surgery (70). Perhaps, studies should focus first on providing definitive proof for surgical treatment of multiple simple rib fractures. The upcoming FixCon trial could possibly provide more exclusive evidence on the clinical, functional, and economic outcomes of operative fixation versus non-operative treatment of patients with multiple simple rib fractures (71).

Future perspectives on management

The current potential of peripheral nerve blocks is believed to surpass neuraxial techniques in the future (20). Although the concept of cryoanalgesia dates back to 1962 (72), this peripheral nerve block technique has gained wide popularity for pain control in thoracic surgery over the past years. In SSRF, cryoablation of the intercostal nerves has proven an outstanding pain management modality, premised on the cost-effectiveness by reducing length of hospital and ICU stay, and decreased narcotic consumption compared to SSRF with conventional pain management protocols (73,74). Still, the addition of cryoanalgesia to SSRF may feel as an overshoot since one of the main indications for SSRF is pain control through fracture stabilization. With the option of percutaneous cryoablation it would be interesting to compare cryoanalgesia as stand-alone nonoperative treatment versus SSRF among patients with stable unilateral rib fractures and high pain scores indicating surgical treatment. Thus, future research should focus on elucidating the efficacy of stand-alone cryoanalgesia for rib fracture pain management.

Numerous factors attribute to the success of SSRF, with an adequate reconstruction of the chest wall being key, allowing for proper expansion and retraction of the rib cage during breathing. Due to the complex shape of the ribs in the chest wall, it remains challenging for fixation devices to fully conform to the anatomical structure of the fractured rib. Currently, several studies have been reported using pre-operative 3D printing derived from CT scan data, offering more insight from all angles into the fracture itself as well as the specific anatomical structure of the rib (55,75,76). Hence, these 3D printing molds can be used for pre-bending of the plates and tailoring of the fixation device, saving operative time, and reducing fixation failure in the future. However, to date it remains unknown whether the addition of 3D printing translates into improved patient outcome, and whether it is cost-effective.

Conclusions

Rib fractures after traumatic thoracic injury can effectively be managed by a patient-tailored non-operative multimodal pain strategy as well as surgical stabilization for selected patients. In both treatment options, it is important to consider a multidisciplinary solution in which pain relief is the key priority. Focusing less on a one-size-fits-all solution and more on future developments, such as percutaneous cryoablation of the intercostal nerves as part of conservative management, as well as the application of 3D printing and use of bio-absorbable materials in the surgical treatment of rib fractures, could possibly benefit patients with rib fractures the most.

Acknowledgments

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned by the editorial office, *Journal of Thoracic Disease*, for the series "Chest Wall Resections and Reconstructions". The article has undergone external peer review.

Peer Review File: Available at https://jtd.amegroups.com/article/view/10.21037/jtd-23-1832/prf

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at https://jtd.amegroups.com/article/view/10.21037/jtd-23-1832/coif). The special series "Chest Wall Resections and Reconstructions" was commissioned by the editorial office without any funding or sponsorship. J.H.T.D. and E.R.d.L. served as the unpaid

© Journal of Thoracic Disease. All rights reserved.

Guest Editors of the series. The authors have no other conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work, ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: https://creativecommons.org/licenses/by-nc-nd/4.0/.

References

1. Nathens A, Fantus R. National Trauma Data Bank 2010 Annual Report. Chicago: American College Surgeon; 2010.
2. Peek J, Beks RB, Hietbrink F, et al. Epidemiology and outcome of rib fractures: a nationwide study in the Netherlands. Eur J Trauma Emerg Surg 2022;48:265-71.
3. ATLS Subcommittee; American College of Surgeons' Committee on Trauma; International ATLS Working Group. Advanced trauma life support (ATLS®): the ninth edition. J Trauma Acute Care Surg 2013;74:1363-6.
4. Vafaei A, Hatamabadi HR, Heidary K, et al. Diagnostic Accuracy of Ultrasonography and Radiography in Initial Evaluation of Chest Trauma Patients. Emerg (Tehran) 2016;4:29-33.
5. Tataroglu O, Erdogan ST, Erdogan MO, et al. Diagnostic Accuracy of InitiaI Chest X-Rays in Thorax Trauma. J Coll Physicians Surg Pak 2018;28:546-8.
6. Livingston DH, Shogan B, John P, et al. CT diagnosis of Rib fractures and the prediction of acute respiratory failure. J Trauma 2008;64:905-11.
7. Santorelli JE, Chau H, Godat L, et al. Not so FAST-Chest ultrasound underdiagnoses traumatic pneumothorax. J Trauma Acute Care Surg 2022;92:44-8.
8. Hwang EG, Lee Y. Simple X-ray versus ultrasonography examination in blunt chest trauma: effective tools of accurate diagnosis and considerations for rib fractures. J Exerc Rehabil 2016;12:637-41.
9. Expert Panel on Thoracic Imaging; Henry TS, Donnelly EF, et al. ACR Appropriateness Criteria(®) Rib Fractures. J Am Coll Radiol 2019;16:S227-34.
10. Griffith JF, Rainer TH, Ching AS, et al. Sonography compared with radiography in revealing acute rib fracture. AJR Am J Roentgenol 1999;173:1603-9.
11. Hurley ME, Keye GD, Hamilton S. Is ultrasound really helpful in the detection of rib fractures? Injury 2004;35:562-6.
12. LaBan MM, Siegel CB, Schutz LK, et al. Occult radiographic fractures of the chest wall identified by nuclear scan imaging: report of seven cases. Arch Phys Med Rehabil 1994;75:353-4.
13. Flagel BT, Luchette FA, Reed RL, et al. Half-a-dozen ribs: the breakpoint for mortality. Surgery 2005;138:717-23; discussion 723-5.
14. Peek J, Smeeing DPJ, Hietbrink F, et al. Comparison of analgesic interventions for traumatic rib fractures: a systematic review and meta-analysis. Eur J Trauma Emerg Surg 2019;45:597-622.
15. Jensen CD, Stark JT, Jacobson LL, et al. Improved Outcomes Associated with the Liberal Use of Thoracic Epidural Analgesia in Patients with Rib Fractures. Pain Med 2017;18:1787-94.
16. Simon B, Ebert J, Bokhari F, et al. Management of pulmonary contusion and flail chest: an Eastern Association for the Surgery of Trauma practice management guideline. J Trauma Acute Care Surg 2012;73:S351-61.
17. Elawamy A, Morsy MR, Ahmed MAY. Comparison of Thoracic Erector Spinae Plane Block With Thoracic Paravertebral Block for Pain Management in Patients With Unilateral Multiple Fractured Ribs. Pain Physician 2022;25:483-90.
18. Bachoumas K, Levrat A, Le Thuaut A, et al. Epidural analgesia in ICU chest trauma patients with fractured ribs: retrospective study of pain control and intubation requirements. Ann Intensive Care 2020;10:116.
19. Duch P, Møller MH. Epidural analgesia in patients with traumatic rib fractures: a systematic review of randomised controlled trials. Acta Anaesthesiol Scand 2015;59:698-709.
20. Koushik SS, Bui A, Slinchenkova K, et al. Analgesic Techniques for Rib Fractures-A Comprehensive Review Article. Curr Pain Headache Rep 2023;27:747-55.
21. Rosero EB, Joshi GP. Nationwide incidence of serious complications of epidural analgesia in the United States. Acta Anaesthesiol Scand 2016;60:810-20.
22. Hardin KS, Leasia KN, Haenel J, et al. The Sequential Clinical Assessment of Respiratory Function (SCARF) score: A dynamic pulmonary physiologic score that

© Journal of Thoracic Disease. All rights reserved.

predicts adverse outcomes in critically ill rib fracture patients. J Trauma Acute Care Surg 2019;87:1260-8.
23. Battle C, Hutchings H, Lovett S, et al. Predicting outcomes after blunt chest wall trauma: development and external validation of a new prognostic model. Crit Care 2014;18:R98.
24. Seok J, Cho HM, Kim HH, et al. Chest Trauma Scoring Systems for Predicting Respiratory Complications in Isolated Rib Fracture. J Surg Res 2019;244:84-90.
25. van Gool MH, van Roozendaal LM, Vissers YLJ, et al. VATS-assisted surgical stabilization of rib fractures in flail chest: 1-year follow-up of 105 cases. Gen Thorac Cardiovasc Surg 2022;70:985-92.
26. Cataneo AJ, Cataneo DC, de Oliveira FH, et al. Surgical versus nonsurgical interventions for flail chest. Cochrane Database Syst Rev 2015;2015:CD009919.
27. Spronk I, Van Wijck SFM, Van Lieshout EMM, et al. Rib Fixation for Multiple Rib Fractures: Healthcare Professionals Perceived Barriers and Facilitators to Clinical Implementation. World J Surg 2023;47:1692-703.
28. Choi J, Khan S, Sheira D, et al. Prospective study of long-term quality-of-life after rib fractures. Surgery 2022;172:404-9.
29. Prins JTH, Van Lieshout EMM, Overtoom HCG, et al. Long-term pulmonary function, thoracic pain, and quality of life in patients with one or more rib fractures. J Trauma Acute Care Surg 2021;91:923-31.
30. Pieracci FM, Majercik S, Ali-Osman F, et al. Consensus statement: Surgical stabilization of rib fractures rib fracture colloquium clinical practice guidelines. Injury 2017;48:307-21.
31. Delaplain PT, Schubl SD, Pieracci FM, et al. Chest wall injury society guideline for SSRF indications, contraindications and timing. Chest Wall Injury Society; 2020. Available online: https://cwisociety.org/wp-content/uploads/2020/05/CWIS-SSRF-Guideline-01102020.pdf
32. Freitag P, Bechmann C, Eden L, et al. Surgical stabilization of serial rib fractures is advantageous in patients with relevant traumatic brain injury. Eur J Trauma Emerg Surg 2022;48:3237-42.
33. Van Wijck SFM, Pieracci FM, Smith EF, et al. Rib fixation in patients with severe rib fractures and pulmonary contusions: Is it safe? J Trauma Acute Care Surg 2022;93:721-6.
34. Prins JTH, Van Lieshout EMM, Ali-Osman F, et al. Outcome after surgical stabilization of rib fractures versus nonoperative treatment in patients with multiple rib fractures and moderate to severe traumatic brain injury (CWIS-TBI). J Trauma Acute Care Surg 2021;90:492-500.
35. Prins JTH, Wijffels MME, Pieracci FM. What is the optimal timing to perform surgical stabilization of rib fractures? J Thorac Dis 2021;13:S13-25.
36. Choi J, Tennakoon L, You JG, et al. Pulmonary contusions in patients with rib fractures: The need to better classify a common injury. Am J Surg 2021;221:211-5.
37. Lagazzi E, Rafaqat W, Argandykov D, et al. Timing matters: Early versus late rib fixation in patients with multiple rib fractures and pulmonary contusion. Surgery 2024;175:529-35.
38. Schots JP, Vissers YL, Hulsewé KW, et al. Addition of Video-Assisted Thoracoscopic Surgery to the Treatment of Flail Chest. Ann Thorac Surg 2017;103:940-4.
39. Bae CM, Son SA, Lee YJ, et al. Clinical Outcomes of Minimally Invasive Surgical Stabilization of Rib Fractures Using Video-Assisted Thoracoscopic Surgery. J Chest Surg 2023;56:120-5.
40. Bui JT, Browder SE, Wilson HK, et al. Does routine uniportal thoracoscopy during rib fixation identify more injuries and impact outcomes? J Thorac Dis 2020;12:5281-8.
41. Mischler D, Schopper C, Gasparri M, et al. Is intrathoracic rib plate fixation advantageous over extrathoracic plating? A biomechanical cadaveric study. J Trauma Acute Care Surg 2022;92:574-80.
42. Tichenor M, Reparaz LB, Watson C, et al. Intrathoracic plates versus extrathoracic plates: a comparison of postoperative pain in surgical stabilization of rib fracture technique. Trauma Surg Acute Care Open 2023;8:e001201.
43. Pieracci FM, Johnson JL, Stovall RT, et al. Completely thoracoscopic, intra-pleural reduction and fixation of severe rib fractures. Trauma Case Rep 2015;1:39-43.
44. Yang Z, Wen M, Kong W, et al. Complete uni-port video-assisted thoracoscopic surgery for surgical stabilization of rib fractures: a case report. J Cardiothorac Surg 2023;18:61.
45. Fokin AA, Hus N, Wycech J, et al. Surgical Stabilization of Rib Fractures: Indications, Techniques, and Pitfalls. JBJS Essent Surg Tech 2020;10:e0032.
46. Bottlang M, Walleser S, Noll M, et al. Biomechanical rationale and evaluation of an implant system for rib fracture fixation. Eur J Trauma Emerg Surg 2010;36:417-26.
47. Hojski A, Xhambazi A, Wiese MN, et al. Chest wall stabilization and rib fixation using a nitinol screwless system in selected patients after blunt trauma: long-term results in a single-centre experience. Interact Cardiovasc

© Journal of Thoracic Disease. All rights reserved.

Thorac Surg 2022;34:386-92.
48. Choke A, Wong YR, Joethy JV. Biomechanical comparison of monocortical and bicortical plate fixation for rib fractures in a cadaveric model using a locking plate system. J Thorac Dis 2019;11:4966-71.
49. Divisi D, Mucilli F, Leonardo GD, et al. Plates versus struts versus an extracortical rib fixation in flail chest patients: Two-center experience. Injury 2021;52:235-42.
50. Puma F, Ragusa M, Santoprete S, et al. As originally published in 1992: Chest wall stabilization with synthetic reabsorbable material. Updated in 1999. Ann Thorac Surg 1999;67:1823-4.
51. Länsman S, Serlo W, Linna O, et al. Treatment of pectus excavatum with bioabsorbable polylactide plates: Preliminary results. J Pediatr Surg 2002;37:1281-6.
52. Mayberry JC, Terhes JT, Ellis TJ, et al. Absorbable plates for rib fracture repair: preliminary experience. J Trauma 2003;55:835-9.
53. Oyamatsu H, Ohata N, Narita K. New technique for fixing rib fracture with bioabsorbable plate. Asian Cardiovasc Thorac Ann 2016;24:736-8.
54. Marasco SF, Sutalo ID, Bui AV. Mode of failure of rib fixation with absorbable plates: a clinical and numerical modeling study. J Trauma 2010;68:1225-33.
55. Waseda R, Matsumoto I, Tatsuzawa Y, et al. Successful Surgical Fixation Using Bio-Absorbable Plates for Frail Chest in a Severe Osteoporotic Octogenarian. Ann Thorac Cardiovasc Surg 2019;25:336-9.
56. Peek J, Beks RB, Hietbrink F, et al. Complications and outcome after rib fracture fixation: A systematic review. J Trauma Acute Care Surg 2020;89:411-8.
57. Mvoula L, Skubic J, Weaver D, et al. Morbidity and Mortality After Rib Fracture in Elderly Patients (>65 Years Old) Compared to a Younger Cohort (≤65 Years of Age) at Doctor Hospital Renaissance Health. Cureus 2022;14:e30941.
58. Pressley CM, Fry WR, Philp AS, et al. Predicting outcome of patients with chest wall injury. Am J Surg 2012;204:910-3; discussion 913-4.
59. He W, Yang Y, Salonga R, et al. Surgical stabilization of multiple rib fractures in an Asian population: a systematic review and meta-analysis. J Thorac Dis 2023;15:4961-75.
60. Niziolek G, Goodman MD, Makley A, et al. "Early results after initiation of a rib fixation programme: A propensity score matched analysis". Injury 2022;53:137-44.
61. Shiroff AM, Wolf S, Wu A, et al. Outcomes of surgical versus nonsurgical treatment for multiple rib fractures: A US hospital matched cohort database analysis. J Trauma Acute Care Surg 2023;94:538-45.
62. Tran Z, Cho NY, De Polo N, et al. Association of Early Rib Plating on Clinical and Financial Outcomes: A National Analysis. Am Surg 2024;90:754-61.
63. Uchida K, Miyashita M, Kaga S, et al. Long-term outcomes of surgical rib fixation in patients with flail chest and multiple rib fractures. Trauma Surg Acute Care Open 2020;5:e000546.
64. Majercik S, Cannon Q, Granger SR, et al. Long-term patient outcomes after surgical stabilization of rib fractures. Am J Surg 2014;208:88-92.
65. Peek J, Kremo V, Beks R, et al. Long-term quality of life and functional outcome after rib fracture fixation. Eur J Trauma Emerg Surg 2022;48:255-64.
66. Pieracci FM, Leasia K, Hernandez MC, et al. Surgical stabilization of rib fractures in octogenarians and beyond-what are the outcomes? J Trauma Acute Care Surg 2021;90:1014-21.
67. Granetzny A, Abd El-Aal M, Emam E, et al. Surgical versus conservative treatment of flail chest. Evaluation of the pulmonary status. Interact Cardiovasc Thorac Surg 2005;4:583-7.
68. Liu T, Liu P, Chen J, et al. A Randomized Controlled Trial of Surgical Rib Fixation in Polytrauma Patients With Flail Chest. J Surg Res 2019;242:223-30.
69. Marasco SF, Davies AR, Cooper J, et al. Prospective randomized controlled trial of operative rib fixation in traumatic flail chest. J Am Coll Surg 2013;216:924-32.
70. Meyer DE, Harvin JA, Vincent L, et al. Randomized Controlled Trial of Surgical Rib Fixation to Nonoperative Management in Severe Chest Wall Injury. Ann Surg 2023;278:357-65.
71. Wijffels MME, Prins JTH, Polinder S, et al. Early fixation versus conservative therapy of multiple, simple rib fractures (FixCon): protocol for a multicenter randomized controlled trial. World J Emerg Surg 2019;14:38.
72. Cooper SM, Dawber RP. The history of cryosurgery. J R Soc Med 2001;94:196-201.
73. Bauman ZM, Loftus J, Raposo-Hadley A, et al. Surgical stabilization of rib fractures combined with intercostal nerve cryoablation proves to be more cost effective by reducing hospital length of stay and narcotics. Injury 2021;52:1128-32.
74. Marturano MN, Thakkar V, Wang H, et al. Intercostal nerve cryoablation during surgical stabilization of rib fractures decreases post-operative opioid use, ventilation days, and intensive care days. Injury 2023;54:110803.
75. Bergquist JR, Morris JM, Matsumoto JM, et al. 3D printed

© Journal of Thoracic Disease. All rights reserved.

modeling contributes to reconstruction of complex chest wall instability. Trauma Case Rep 2019;22:100218.
76. Zhou XT, Zhang DS, Yang Y, et al. Analysis of the advantages of 3D printing in the surgical treatment of multiple rib fractures: 5 cases report. J Cardiothorac Surg 2019;14:105.

Cite this article as: Franssen AJPM, Daemen JHT, Luyten JA, Meesters B, Pijnenburg AM, Reisinger KW, van Vugt R, Hulsewé KWE, Vissers YLJ, de Loos ER. Treatment of traumatic rib fractures: an overview of current evidence and future perspectives. J Thorac Dis 2024;16(8):5399-5408. doi: 10.21037/jtd-23-1832

© Journal of Thoracic Disease. All rights reserved.